



Heart Disease

Heart Disease

Risk Factors for Coronary Heart Disease

The American Heart Association has identified several risk factors that contribute to Coronary Heart Disease. Some of them can be modified, treated or controlled, and some can't. The more risk factors you have, the greater your chance of developing coronary heart disease. Also, the greater the level of each risk factor, the greater the risk. For example, a person with a total cholesterol of 300 mg/dL, even though everyone with a total cholesterol greater than 240 is considered high-risk.

What Risk Factors Can't Be Changed?

Increasing Age - Over 83% of people who die of coronary heart disease are 65 or older. At older ages, women who have heart attacks are more likely than men are to die from them within a few weeks.

Male Sex (Gender) - Men have a greater risk of heart attack than women do, and they have attacks earlier in life. Even after menopause, when women's death rate from heart disease increases, it's not as great as men's.

Heredity (Including Race) - Children of parents with heart disease are more likely to develop it themselves. African Americans have more severe high blood pressure than Caucasians and a higher risk of heart disease. Heart Disease risk is also higher among Mexican Americans, American Indians, native Hawaiians and some Asian Americans. This is partly due to higher rates of obesity and diabetes have one or more other risk factors. Just as you can't control your age, sex and race, you can't control your family history. Therefore, it's even more important to treat and control any other risk factors you have.

What Risk Factors Can You Modify?

Tobacco Smoke - Smokers' risk of developing coronary heart disease is 2-4 times that of nonsmokers. Cigarette smoking is a powerful independent risk factor for sudden cardiac death in patients with coronary heart disease; smokers have about twice the risk of nonsmokers. People who smoke cigars or pipes seem to have a higher risk of death from coronary heart disease (and possibly stroke) but their risk isn't as great as cigarette smokers'. Exposure to other people's smoke increases the risk of heart disease even for nonsmokers.

High Blood Cholesterol - As blood cholesterol rises, so does the risk of coronary heart disease. When other risk factors are present, this risk increases even more. A person's cholesterol level is also affected by age, sex, heredity and diet.

High Blood Pressure - High blood pressure increases the heart's workload, causing the heart to thicken and become stiffer. It also increases your risk of stroke, heart attack, kidney failure and congestive heart failure.

Physical Inactivity - An inactive lifestyle is a risk factor for coronary heart disease. Regular, moderate-to-vigorous physical activity helps prevent heart and blood vessel disease. The more vigorous the activity, the greater your benefits. However, even more moderate-intensity activities help if done regularly and long term.

Obesity and Overweight - People who have excess body fat - especially if a lot of it is at the waist - are more likely to develop heart disease and stroke even if they have no other risk factors. Excess weight increases the heart's work. It also raises blood pressure and blood cholesterol and triglyceride levels, and lowers HDL ("good") cholesterol levels. It can also make diabetes more likely to develop.

Diabetes Mellitus - Diabetes seriously increases your risk of developing cardiovascular disease. Even when glucose (blood sugar) levels are under control, diabetes increases the risk of heart disease and stroke, but the risks are even greater if blood sugar is not well controlled.

STAY CONNECTED



Wellness information courtesy of Wellness Proposals